

PATENT
Attorney Docket No. AMBER-06803

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Kenneth J. Rothschild *et al.*

Serial No.: 10/034,736

Group No.: 1636


Filed: 12/27/01

Examiner: Ketter, J.

Entitled: **Detection Of Markers In Nascent Proteins**

INFORMATION DISCLOSURE STATEMENT

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)	
I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.	
Dated: <u>March 30, 2004</u>	By:  Christopher J. Collins

Sir:

The citations listed below, copies will follow under separate cover, may be material to the examination of the above-identified application, and are therefore submitted in compliance with the duty of disclosure defined in 37 C.F.R. §§ 1.56 and 1.97. The Examiner is requested to make these citations of official record in this application.

The following printed publications are referred to in the body of the specification:

- U. S. Patent No. 5,190,632 to Fujimiya *et al.*;
- U. S. Patent No. 5,069,769 to Fujimiya *et al.*;
- U. S. Patent No. 5,137,609 to Manian *et al.*;
- U. S. Patent No. 4,683,195 to Mullis *et al.*;
- U. S. Patent No. 4,241,050 to Barth;
- U. S. Patent No. 4,728,591 to Clark *et al.*;
- U. S. Patent No. 4,802,951 to Clark *et al.*;
- PCT WO90/05785 to Shultz;
- Noren, *et al.*, "A General Method for Site-Specific Incorporation of Unnatural Amino Acids into Proteins," *Science* 244:182-188 (1989);

- Bain, *et al.*, "Site-Specific Incorporation of Nonnatural Residues during In Vitro Protein Biosynthesis with Semisynthetic Aminoacyl-tRNAs," *Biochemistry* 30:5411-21 (1991);
- Promega Technical Bulletin No. 182; tRNA^{nascent}[™]: Non-Radioactive Translation Detection System. 1-16, Sept. 1993;
- Krieg, *et al.*, Photocrosslinking of the signal sequence of nascent preprolactin to the 54-kilodalton polypeptide of the signal recognition particle," *Proc. Natl. Acad. Sci. USA* 83:8604-08 (1986);
- Molecular Cell Biology, J. Darnell *et al.*, Editors. Scientific American Books. N.Y., N.Y. 1991;
- *Current Protocols in Molecular Biology* (F.M. Ausubel *et al.* editors, Wiley Interscience, 1993), pp. 10-16, 10-77;
- Seong and RajBhandary, "Escherichia coli formylmethionine tRNA: Mutations in GGG sequence conserved in anticodon stem of initiator tRNAs affect initiation of protein synthesis and conformation of anticodon loop," *Proc. Natl. Acad. Sci. USA* 84:334-338 (1987);
- Bruce and Uhlenbeck, "Specific Interaction of Anticodon Loop Residues with Yeast Phenylalanyl-tRNA Synthetase," *Biochemistry* 21:3921-3926 (1982);
- Pratt, "Coupled Transcription-Translation in Prokaryotic Cell-Free System," (*Transcription and Translation*, B.D. Hames and S.J. Higgins, Editors, p. 179-209, IRL Press, Oxford, 1984);
- Spirin, *et al.*, "A Continuous Cell-Free Translation System Capable of Producing Polypeptides in High Yield," *Sci.* 242:1162-64 (1988);
- Felgner *et al.*, "Lipofection: A highly efficient, lipid-mediated DNA-transfection procedure," *Proc. Natl. Acad. Sci. USA* 84:7413-17 (1987);
- Neu and Heppel, "Nucleotide Sequence Analysis of Polyribonucleotides by Means of Periodate Oxidation Followed by Cleavage with an Amine," *J. Biol. Chem.* 239:2927-34 (1964);
- Sampson and Uhlenbeck, "Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed *in vitro*," *Proc. Natl. Acad. Sci. USA* 85:1033-37 (1988);

- Hudson, "Methodological Implications of Simultaneous Solid-Phases Peptide Synthesis. 1. Comparison of Different Coupling Procedures," *J. Org. Chem.* 53:617-624 (1988);
- Happ, *et al.*, "New Approach to the Synthesis of 2'(3')-O-Aminoacyl Oligoribonucleotides," *J. Org. Chem.* 52:5387-91 (1987);
- Heckler *et al.*, "T4 RNA Ligase Mediated Preparation of Novel "Chemically Misacylated" tRNA ^{Phe}s," *Biochemistry* 23:1468-73 (1984);
- Heckler *et al.*, "Preparation of 2'(3')-O-Acyl-pCpA Derivatives as Substrates for T4 RNA Ligase-Mediated "Chemical Aminoacylation"," *Tetrahedron* 40:87-94 (1984);
- DiCesare, *et al.*, "A High-Sensitivity Electrochemiluminescence-Based Detection System for Automated PCR Product Quantitation," *BioTechniques* 15:152-59 (1993);
- Allen, *et al.*, *Gel Electrophoresis and Isoelectric Focusing of Proteins*, Walter de Gruyter, New York 1984, pp. 17-62;
- Stephen, "High-Resolution Preparative SDS-Polyacrylamide Gel Electrophoresis: Fluorescent Visualization and Electrophoretic Elution-Concentration of Protein Bands," *Anal. Biochem.* 65:369-79 (1975);
- Pillai, "Photoremovable Protecting Groups in Organic Synthesis," *Synthesis* 1-26 (1980);
- Patchornik, *et al.*, "Photosensitive Protecting Groups," *J. Am. Chem. Soc.* 92:6333-35 (1970);
- *Antibodies: A Laboratory Manual* (E. Harlow and D. Lane, editors, Cold Spring Harbor Laboratory Press), 1988, pp. 53,72-73;
- Powell, *et al.*, "Molecular Diagnosis of Familial Adenomatous Polyposis," *N. Engl. J. Med.* 329:1982-87 (1993);
- Sonar, *et al.*, *Biochem.* 32:13777-81, 1993¹;
- Widder, *et al.*, "Magnetic Microspheres: A Model System for Site Specific Drug Delivery in *Vivo*," *Proc. Soc. Exp. Biol. & Med.* 58:141-46 (1978); and

¹ We have been unable to obtain this reference. If the examiner so requests, we can increase our efforts to obtain a copy.

- Oesterhelt *et al.*, "Bacteriorhodopsin: a biological material for information processing," *Quart. Rev. Biophys.* 4:425-78 (1991).

Applicants have become aware of the following printed publications which may be material to the examination of this application:

- US 5,614,386 to Metzker, *et al.*, reports the use of BODIPY fluorophores for labeling DNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- US 5,215,927 to Berenson, *et al.*, describes a method of immunoselection of cells based on avidin and biotinylated antibody. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- US 5,434,272 to Corrie, *et al.*, describes the synthesis of caged fluorophores based on the dyes fluorescein and rhodamine. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- European Pat. No. 0234799A2 to Kurzchalia, *et al.*, describes methods for the detection and isolation of protein utilizing the incorporation of photoaffinity reagents and biotin or other haptens into the lysine residues of nascent peptides. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and the binding of the nascent protein to a surface.
- U.S. Pat. No. 4,675,285 to Clark, *et al.*, describes a method for identification of clones expressing the desired protein from the cDNA libraries. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Sonar, S., *et al.*, "Cell-Free Synthesis, Functional Refolding and Spectroscopic Characteristics of Bacteriorhodopsin, an Integral Membrane Protein," *Biochemistry* 32:13777-13781 (1993) report on the cell-free synthesis of the protein bacteriorhodopsin. The publication does not disclose the

misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Karolin, J., *et al.*, "Flourescence and Absorption Spectroscopic Properties of Dipyrrometheneboron Difluoride (BODIPY) Derivatives in Liquids, lipid Membranes and Proteins," *J. Am. Chem. Soc.* 116:7801-7806 (1994) report on the spectroscopic properties of BODIPY in proteins. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Kurzchalia, T.V., *et al.*, "tRNA-Mediated Labelling of Proteins with Biotin. A Nonradioactive Method for the Detection of Cell-Free Translation Products," *Eur. J. Biochem.* 172:663-668 (1988) report on cell-free translation of proteins made with a modified lysyl-tRNA and biotin label. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Baldini, *et al.*, "Mischarging *Escherichia coli* tRNA^{Phe} with L-4'-[3-Trifluoromethyl)-3*H*-diazirin-3-yl]phenylalanine, a Photoactivatable Analogue of Phenylalanine," *Biochemistry* 27:7951-7959 (1988) report a procedure for the misaminoacylation of tRNA with a photoactivatable analog of phenylalanine. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Hall, *et al.*, "Mapping Labeled Sites in *Escherichia coli* Ribosomal RNA: Distribution of Methyl Groups and Identification of a Photoaffinity-Labeled RNA Region Putatively at the Peptidyltransferase Center," *Biochemistry* 24:5702-5711 (1985) report a method utilizing radiolabeled 5-Azido-2-nitrobenzoyl-[H³]Phe-tRNA^{Phe} to determine structural characteristics of the rRNA in ribosomes. [H³] was detected by electrophoresis of the rRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Billington, *et al.*, "Synthesis and Photochemistry of Photolabile *N*-Glycine Derivatives and Efforts of One on the Glycine Receptor," *Biochemistry* 31:5500-5507 (1992) report the synthesis of photolabile derivatives of *N*-

glycine. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Mouton, *et al.*, "A Reagent for Covalently Attaching Biotin to Proteins via a Cleavable Connector Arm," *Archives of Biochemistry and Biophysics* 218(1):101-108 (1982) report procedures for the attachment of biotin to proteins via a cleavable connector arm. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Thiele and Fahernholz, "Photocleavable Biotinylated Ligands for Affinity Chromatography," *Analytical Biochemistry* 218:330-337 (1994) report a procedure utilizing biotinylated ligands for affinity chromatography. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Herman, *et al.*, "Affinity Chromatography of DNA Labeled with Chemically Cleavable Biotinylated Nucleotide Analogs," *Analytical Biochemistry* 156:48-55 (1986) report methods for the affinity chromatography of DNA labeled with chemically cleavable biotinylated nucleotide analogs. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Perri, *et al.*, "Tandem Photochemical Synthesis of N-Amino β -Lactams from Pyrazolidin-3-ones," *J. Org. Chem.* 55:6037-6047 (1990) report the photochemical synthesis of N-amino beta-lactams from pyrazolidin-3-ones. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Amit, *et al.*, "Photosensitive Protecting Groups of Amino Sugars and Their Use in Glycoside Synthesis. 2-Nitrobenzyloxycarbonylamino and 6-Nitroveratryloxycarbonylamino Derivatives," *J. Org. Chem.* 39(2):192-196 (1974) report the use of 2-nitrobenzyloxycarbonyl and 6-nitroveratryloxycarbonyl as protecting groups for the amino function in amino sugars. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Zehavi, *et al.*, "Light-Sensitive Glycosides. I. 6-Nitroveratryl β -D-Glucopyranoside and 2-Nitrobenzyl β -D-Glucopyranoside," *J. Org. Chem.* 37(14):2281-2285 (1972) report the production and characterization of two light sensitive glucosides, 2-nitrobenzyl beta-D-glucopyranoside and 6-nitroveratryl. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Peyser and Flechtner, "*N*-(α -Hydroxy-2-nitrosobenzyl)-1-naphthamide: A Photochemical Intermediate," *J. Org. Chem.* 25:4645-4646 (1987) report the synthesis of an unstable nitro compound intermediate resulting from the irradiation of *N*-(2-nitrobenzyl)-1-naphthamide. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Ohtsuka, *et al.*, "Studies on Transfer Ribonucleic Acids and Related Compounds. 20. A New Versatile Ribooligonucleotide Block with 2'-(*o*-Nitrobenzyl) and 3'-Phosphorodianilidate Groups Suitable for Elongation of Chains in the 3' and 5' Directions," *J. Am. Chem. Soc.* 100(14):4580-4584 (1978) report methods to protect the 2'-hydroxyl group of ribonucleotides during the synthesis of ribonucleotides. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Mendel, *et al.*, "Construction of a Light-Activated Protein by Unnatural Amino Acid Mutagenesis," *J. Am. Chem. Soc.* 113:2758-2760 (1991) report the construction of a tRNA for use in the *in vitro* translation synthesis of a caged T4L lysozyme. The modified lysozyme is inactive until irradiated where upon the enzyme gains full activity. The modified, activated lysozyme is detected by its ability to lyse *E. coli* by donating a proton to the interglycosidic oxygen of beta(1 \rightarrow 4)-linked NAM-NAG residues in the cell wall thereby giving a visual readout. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.

- Robertson, *et al.*, "A General and Efficient Route for Chemical Aminoacylation of Transfer RNAs," *J. Am. Chem. Soc.* 113:2722-2729 (1991) report a method for the misaminoacylation of tRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Miknis and Williams, "Total Synthesis of (\pm)-Aspirochlorine," *J. Am. Chem. Soc.* 115:536-547 (1993) report the synthesis of (\pm)-aspirochlorine which utilized a 2-nitrobenzyl group as an amide protecting group. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Dawson, *et al.*, "Affinity Isolation of Transcriptionally Active Murine Erythroleukemia Cell DNA Using a Cleavable Biotinylated Nucleotide Analog," *J. Biological Chemistry* 264:12830-12837 (1989) report a method for the affinity isolation of transcriptionally active cellular DNA using a biotinylated nucleotide analog. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Markings and Tsien, "Caged Nitric Oxide, Stable organic Molecules from which Nitric Oxide can be Photoreleased," *J. Biological Chemistry* 269:6282-6285 (1994) report the synthesis and testing of a series of caged nitric oxide compounds. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Thompson, *et al.*, "Photocleavable Nitrobenzyl-Protein Conjugates," *Biochemical and Biophysical Research Communications* 201(3):1213-1219 (1994) report on the synthesis of photocleavable nitrobenzyl-protein conjugates. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Albarella, *et al.*, "Monoadduct forming photochemical reagents for labeling nucleic acids for hybridization," *Nucleic Acids Research* 17(11):4293-4308 (1989) report the synthesis of novel photoreagents for use in the labeling of DNA with biotin. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Robertson, *et al.*, "The use of 5'-phospho-2 deoxyribocytidylylriboadenosine as a facile route to chemical aminoacylation of tRNA," *Nucleic Acids Research* 17(23):9649-9660 (1989) report a method of misaminoacylating tRNA, performing *in vitro* translation with the misaminoacylated tRNA, and detecting the translated protein with incorporation of [³⁵S] Met into the nascent peptide during translation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Lesnikowski and Jaworska, "Studies on Stereospecific Formation of P-Chiral Internucleotide Linkage. Synthesis of (Rp,Rp)- and (Sp,Sp)-Thymidylyl(3',5') Thymidylyl (3',5') Thymidine DI(O,O-Phosphorothioate) using 2-Nitrobenzyl Group as a New S-Protection," *Tetrahedron Letters* 30(29):3821-3824 (1989) report studies on the stereospecific formation of p-chiral internucleotide linkages. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Renil and Pillai, "Synthesis of fully Protected Peptides on a Tetraethyleneglycol Diacrylate (TTEGDA)-Crosslinked Polystyrene support with a Photolytically Detachable 2-Nitrobenzyl Anchoring group," *Tetrahedron Letters* 35(22):3809-3812 (1994) report the synthesis of full peptides on solid support that are released by photolytic cleavage. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Zehavi and Herchman, "Enzymic synthesis of oligosaccharides on a polymer support, light-sensitive, water-soluble substituted poly(vinyl alcohol)," *Carbohydrate Research* 128:160-164 (1984) report an improvement in the synthesis of saccharide derivatives wherein the resulting oligonucleotide is released by irradiation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Whitney, R.A., "A Photochemical approach to the synthesis of (±)-biotin," *Can. J. Chem.* 59:2650-2653 (1981) reports the synthesis of (±)-biotin utilizing

oxidative rearrangement. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Nishikubo, *et al.*, "Study of Photopolymers. XXXIV. Etherification and Esterification Reactions of Polymers with (*o*, *m*, or *p*)-Bromomethylnitrobenzene Using the DBU Method and the Photochemical Properties of the Resulting Polymers," *J. Polymer Science: Part A: Polymer Chemistry* 28:105-117 (1990) report the successful reaction of poly(4-hydroxystyrene) (PHST) and PMAA with *o*-, *m*- or *p*-bromomethylnitrobenzenes using the DBU method in aprotic polar solvents and studies the photochemical properties of the resulting polymers. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Zehavi, *et al.*, "Enzymic Synthesis of Oligosaccharides on a Polymer Support Light-Sensitive, Substituted Polyacrylamide Beads," *Carbohydrate Research* 124:23-34 (1983) report the enzymatic synthesis of two light-sensitive oligosaccharides on a polymer support. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Ohtsuka, *et al.*, "Studies on Transfer Ribonucleic Acids and Related Compounds; XVIII. A Photolabile 2'-Ether of Guanosine as an Intermediate for Oligonucleotide Synthesis," *Synthesis* 7:453-454 (1977) report the characterization of a photolabile 2'-ether of guanosine as an intermediate for oligonucleotide synthesis. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Yen, *et al.*, "Optically controlled ligand delivery, 1: Synthesis of Water-soluble Copolymers Containing Photocleavable Bonds," *Makromol. Chem.* 190:69-82 (1989) report the synthesis of photocleavable polymers for potential use in immunoassays. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Wilchek and Bayer, "Applications of Avidin-Biotin Technology: Literature Survey," *Methods in Enzymology* 184:14-45 (1990) report the use of avidin-biotin technology for protein identification, analysis and purification. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Houlihan, *et al.*, "Nitrobenzyl Ester Chemistry for Polymer Processes Involving Chemical Amplification," *Macromolecules* 21:2001-2006 (1988) report the synthesis of 2-nitrobenzyl, 2,4-dinitrobenzyl and 2,6-dinitrobenzyl tosylate for use as photochemically labile protective groups. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Chemical Analysis, Hemmila, "Application of Fluorescence in Immunoassays," p139-158, reports the physical characteristics and methods of quantification of various organometallic fluorochromes. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Kozak, "Point Mutations Define a Sequence Flanking the AUG Initiator Codon that Modulates Translation by Eukaryotic Ribosomes," *Cell* 44:283-292 (1986) reports a sequence flanking the AUG initiator codon that modulates eukaryotic ribosomes. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Odom, *et al.*, Methods in Molecular Biology "In Vitro Engineering Using Acyl-Derivatized tRNAs," edited by: R. Martin Humana Press Inc., Totowa, NJ Chapter 7, pp.93-103, report the synthesis of CPM-SAc-[³⁵S]Met-tRNA_f by successive reactions of [³⁵S]Met-tRNA with DTDG monosuccinimidyl ester, DTT and CPM. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Varshney, *et al.*, "Initiation of protein synthesis from a termination codon," *PNAS* 87:1586-1590 (1990) report the use of a misaminoacylation of a tRNA to demonstrate that translation can begin with a termination codon. The publication does not disclose the misaminoacylation of tRNA with a non-

radioactive marker followed by a translation step and binding the nascent protein to a surface.

- Varshney, *et al.*, "Direct Analysis of Aminoacylation Levels of tRNAs in Vivo," *J. Biological Chemistry* 266(36):24712-24718 (1991) report methods of detecting aminoacylation levels of tRNAs *in vivo* using mutated tRNAs and gel electrophoresis. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and bonding the nascent protein to a surface.
- Krafft, *et al.*, "Photoactivable Fluorophores. 3. Synthesis and Photoactivation of Fluorogenic Difunctionalized Fluoresceins," *J. Am. Chem. Soc.* 110:301-303 (1988) report the synthesis of a photoactivatable fluorophore designed for tracer studies of molecular transport and diffusion in biological systems. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Kalbag and Roeske, "A Photolabile Protecting Group for Histidine," *J. Am. Chem. Soc.* 97:440-441 (1975) reports the synthesis wherein the *o*-nitrobenzyl group of imidazole was introduced into the side chain of histidine and subsequently removal by irradiation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Crowley, *et al.*, "The signal sequence moves through a ribosomal tunnel into a noncytoplasmic aqueous environment at the ER membrane early in translocation," *Cell* 73:1101-1115 (1993). This reference reports the logistics of nascent peptide production by utilizing [¹⁴C]Lys-tRNA analogs to incorporate fluorescent probes into the signal sequence to examine the environment of a nascent peptide chain as it moved through the ribosome and into the ER membrane. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Hardesty, *et al.*, "Extension and Folding of Nascent Peptides on Ribosomes." The Translational Apparatus, Nierhaus *et al.* ed: New York and London;

Plenum Press. p.347-358 (1993) report peptide folding experiments where amino-tRNAs were modified after aminoacylation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.

- Johnson, *et al.*, "Protein Synthesis and Secretion as seen by the Nascent Protein Chain," The Translational Apparatus, Nierhaus *et al.* ed: New York and London; Plenum Press. p. 359-370 (1993) report peptide synthesis experiments where Lys-tRNA was modified after aminoacylation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Czworkowski, *et al.*, "Fluorescence Study of the Topology of Messenger RNA Bound to the 30S Ribosomal Subunit of *Escherichia coli*," *Biochemistry* 30:4821-4830 (1991), report the interaction of fluorescently labeled RNAs (25-36 nucleotides in length) with the fluorescently labeled 30S subunit of *Escherichia coli* was studied by using fluorescence spectroscopic techniques. By using the distances calculated from energy transfer a topological map of this region of mRNA on the 30S subunit was constructed. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Hardesty, *et al.*, "Ribosome function determined by fluorescence," *Biochimie* 74:391-401 (1992), report on the fluorescence spectroscopy and fluorescent resonance energy transfer and their application to the studies on the structure, movement and conformation of tRNA, the nascent peptide, and mRNA in a ribosome during the reaction steps of peptide elongation. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and and binding the nascent protein to a surface.
- Kudlicki, *et al.*, "Chaperone-dependent Folding and Activation of Ribosome-bound Nascent Rhodanese," *J. Mol. Biol.* 244:319-331 (1994), report on the synthesis of coumarin labeled rhodanese by coupled transcription/translation in a cell-free *Escherichia coli* system. The influence of chaperones on the release of nascent rhodanese from ribosome was studied by fluorescence spectroscopy.

The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Picking, *et al.*, "The use of synthetic tRNA as probes for examining nascent peptides on *Escherichia coli* ribosomes," *Biochimie* 73:1101-1107 (1991), the cell free synthesis of N-acetyl or N-acyl coumarin labeled polycysteine and polyserine was carried out on *Escherichia coli* ribosomes using N-acyl coumarin derivatives of either Ser-tRNA or Phe-tRNA. The properties of the resulting nascent peptides were studied by fluorescence spectroscopy and compared to those of nascent polyphenylalanine chains synthesized under similar conditions. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Picking, *et al.*, "Evidence for RNA in the Peptidyl Transferase Center of *Escherichia coli* Ribosomes as Indicated by Fluorescence," *Biochemistry* 31:12565-12570 (1992). The interaction of coumarin labeled (tRNA(phe)) [either the amino acid or the 5' end] with ribosomes was studied using fluorescence spectroscopy. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Picking, *et al.*, "The Conformation of Nascent Polylysine and Polyphenylalanine Peptides on Ribosomes," *J. Biol. Chem.* 266:1534-1542 (1991), report the behavior of fluorescently labeled polylysine and polyphenylalanine during their *in vitro* synthesis on *E. coli* ribosomes. The position and conformation of the nascent peptide were monitored by fluorescence techniques. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Picking, *et al.*, "Fluorescence Characterization of the Environment Encountered by Nascent Polyalanine and Polyserine as They Exit *Escherichia coli* Ribosomes during Translation," *Biochemistry* 31:2368-2375 (1992) report that a coumarin probe placed at the alpha-amino group of a synthetic elongator

alanyl-tRNA or a synthetic initiator alanyl-tRNA or at the epsilon-amino group of natural lysyl-tRNA, and each was used to nonenzymatically initiate peptide synthesis. The interaction of nascent polypeptides with the environments was studied using fluorescence techniques. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.

- Picking, *et al.*, "A synthetic alanyl-initiator tRNA with initiator tRNA properties as determined by fluorescence measurements: Comparison to a synthetic alanyl-elongator tRNA," *Nucleic Acids Research* 19:5749-5754 (1991) report that a derivative of coumarin was covalently attached to the alpha amino group of alanine of the two synthetic Ala-tRNA species. The fluorescence spectra, quantum yield and anisotropy for the two Ala-tRNA derivatives were studied after they were bound to 70S ribosomes. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Ma, *et al.*, "In Vitro Protein Engineering Using Synthetic tRNA^{Ala} with Different Anticodons," *Biochemistry* 32:7939-7945 (1993), report that synthetic tRNA for *in vitro* protein engineering was tested in a coupled transcription/translation system prepared from *Escherichia coli*. Markers were not attached to the alpha-amino acid. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Odom, *et al.*, "Movement of tRNA but Not the Nascent Peptide during Peptide Bond Formation on Ribosomes," *Biochemistry* 29:10734-10744 (1990), report the interaction of fluorescently labeled (Phe) tRNA (labeled at the 5'-end) with the ribosome as well as with the nascent polypeptide, was investigated using nonradiative energy transfer. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- McCray, "Properties and uses of photoreactive caged compounds," *Annu Rev Biophys Biophys Chem* 18:239-270 (1989), is a review of reports on the


synthesis and uses of caged photoreactive compounds. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.

- Bain, *et al.*, "Site-Specific Incorporation of Nonnatural Residues into Peptides: Effect of Residue Structure on Suppression and Translation Efficiencies" *Tetrahedron* 47:2389-2400 (1991), report on a method for the incorporation of misaminoacylation of tRNA with an amino acid comprising a radioactive tag. Incorporation of the amino acid into a peptide is determined by precipitation and detection of radioactivity by scintillation counting. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Johnson, *et al.*, "N^ε-Acetyllysine Transfer Ribonucleic Acid: A Biologically Active Analogue of Aminoacyl Transfer Ribonucleic Acids" *J Am Chem Soc* 15:569-575 (1976), report a method for the preparation an analogue of Lys-tRNA (ϵ -Ac-Lys-tRNA) for the transfer of the modified amino acid into a nascent peptide chain. Incorporated amino acids were detected by scintillation counter. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Johnson, "Chemically Modified Aminoacyl-tRNA as a Probe of Ribosome Structure: the Synthesis and in vitro Activity of ϵ -N-acetyl-Lys-tRNA," 1973 Thesis Excerpts, University of Oregon, Eugene, Oregon, reports a method for the preparation of fluorescent-labeled Lys-tRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step.
- Shore, S.K., "The use of Fluorescent-Labeled Amino Acids to Examine the Environment of Ribosome-Bound Nascent Polypeptide Chains" Univeristy of Oklahoma, 1991, Thesis, reports a methods for the preparation of fluorescent-labeled Lys-tRNA designated ϵ NBD-Lys-tRNA and ϵ FISAc-Lys-tRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.

- Shore, *et al.*, "A Fluorescent Probe Capable of Incorporation into Nascent Polypeptide Chains," 1986 *Federation Proceedings* 45, 1566 Abstract, report a method for the preparation of fluorescent-labeled Lys-tRNA designated N^ε-bromoacetyl-Lys-tRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.
- Shore, *et al.*, "Accessibility of AA-tRNA and Nascent Chain During Protein Synthesis," 1988 *FASEB Journal* 2, A1045 Abstract, report a method for the preparation of fluorescent-labeled Lys-tRNA designated εFISAc-Lys-tRNA. The publication does not disclose the misaminoacylation of tRNA with a non-radioactive marker followed by a translation step and binding the nascent protein to a surface.

This Information Disclosure Statement under 37 C.F.R. §§ 1.56 and 1.97 is not to be construed as a representation that a search has been made, that additional information material to the examination of this application does not exist, or that any one or more of these citations constitutes prior art.

Dated: March 30, 2004



Peter G. Carroll
Registration No. 32,837

MEDLEN & CARROLL, LLP
101 Howard Street, Suite 350
San Francisco, California 94105
617/984-0616

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: AMBER-06803

Serial No.: 10/034,736

INFORMATION DISCLOSURE STATEMENT BY APPLICANT
(Include Additional Sheets If Necessary)Applicant: Kenneth J. Rothschild *et al.*

Filing Date: 12/27/01

Group Art Unit: 1636

(37 CFR § 1.98(b))

U.S. PATENT DOCUMENTS

Examiner Initials	Cite No.	Serial / Patent Number	Issue Date	Applicant / Patentee	Class	Subclass	Filing Date
	1	4,241,050	12/23/80	Barth	424	114	9/01/78
	2	4,675,285	6/23/87	Clark <i>et al.</i>	435	6	9/19/84
	3	4,683,195	7/28/87	Mullis <i>et al.</i>	435	6	2/07/86
	4	4,728,591	3/01/88	Clark <i>et al.</i>	430	5	3/01/88
	5	4,802,951	2/07/89	Clark <i>et al.</i>	156	630	12/23/87
	6	5,069,769	12/03/91	Fujimiya <i>et al.</i>	204	182.8	6/06/90
	7	5,137,609	8/11/92	Manian <i>et al.</i>	204	180.1	1/31/92
	8	5,190,632	3/02/93	Fujimiya <i>et al.</i>	204	299 R	3/20/92
	9	5,434,272	7/18/95	Corrie <i>et al.</i>	548	525	11/06/91
	10	5,614,386	3/25/97	Metzker <i>et al.</i>	435	91.1	6/23/95

FOREIGN PATENTS OR PUBLISHED FOREIGN PATENT APPLICATIONS

		Document Number	Publication Date	Country / Patent Office	Class	Subclass	Translation	
							Yes	No
	11	EP 0234799 A2	02.09.87	EPO				
	12	WO 90/05785	31.5.90	PCT				

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

	13	Albarella, <i>et al.</i> , "Monoadduct forming photochemical reagents for labeling nucleic acids for hybridization," <i>Nucleic Acids Research</i> 117(11):4293-4308 (1989)
	14	Allen <i>et al.</i> , <i>Gel Electrophoresis and Isoelectric Focusing of Proteins</i> , Walter de Gruyter, New York 1984, pp. 17-62
	15	Amit, <i>et al.</i> , Photosensitive Protecting Groups of Amino Sugars and Their Use in Glycoside Synthesis. 2-Nitrobenzyloxycarbonylamino and 6-Nitroveratryloxycarbonylamino Derivatives," <i>J. Org. Chem.</i> 39(2):192-196 (1974)
	16	<i>Antibodies: A Laboratory Manual</i> (E. Harlow and D. Lane, editors, Cold Spring Harbor Laboratory Press), 1988, pp. 53,72-73
	17	Bain <i>et al.</i> , "Site-Specific Incorporation of Nonnatural Residues during In Vitro Protein Biosynthesis with Semisynthetic Aminoacyl-tRNAs," <i>Biochemistry</i> 30:5411-21 (1991)
	18	Bain <i>et al.</i> , "Site-Specific Incorporation of Non-Natural Residues into Peptides: Effect of Residue Structure on Suppression and Translation Efficiencies," <i>Tetrahedron</i> 47:2389-2400 (1991)
	19	Baldini, <i>et al.</i> , "Mischarging <i>Escherichia coli</i> tRNA ^{Phe} with L-4'-[Trifluoromethyl]-3H-diazirin-3-yl] phenylalanine, a Photoactivatable Analogue of Phenylalanine," <i>Biochemistry</i> 27:7951-7959 (1988)
	20	Billington, <i>et al.</i> , "Synthesis and Photochemistry of Photolabile N-Glycine Derivatives and Efforts of One on the Glycine Receptor," <i>Biochemistry</i> 31:5500-5507 (1992)
	21	Bruce and Uhlenbeck, "Specific Interaction of Anticodon Loop Residues with Yeast Phenylalanyl-tRNA Synthetase," <i>Biochemistry</i> 21:3921-3926 (1982)
	22	Crowley <i>et al.</i> , "The signal sequence moves through a ribosomal tunnel into a noncytoplasmic aqueous environment at the ER membrane early in translocation," <i>Cell</i> 73:1101-1115 (1993)
	23	<i>Current Protocols in Molecular Biology</i> (F.M. Ausubel <i>et al.</i> editors, Wiley Interscience, 1993), pp. 10-16, 10-77
	24	Czworkowski <i>et al.</i> , "Fluorescence Study of the Topology of Messenger RNA Bound to the 30S Ribosomal Subunit of <i>Escherichia coli</i> ," <i>Biochemistry</i> 30:4821-4830 (1991)
	25	Dawson, <i>et al.</i> , "Affinity Isolation of Transcriptionally Active Murine Erythroleukemia Cell DNA Using a Cleavable Biotinylated Nucleotide Analog," <i>J. Biological Chemistry</i> 264:12830-12837 (1989)

Examiner:

Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: AMBER-06803

Serial No.: 10/034,736

INFORMATION MADE AVAILABLE BY APPLICANT

(Use Several Sheets If Necessary)

Applicant: Kenneth J. Rothschild *et al.*

Filing Date: 12/27/01

Group Art Unit: 1636

(37 CFR § 1.98(b))


OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

- | | |
|----|---|
| 26 | DiCesare <i>et al.</i> , "A High-Sensitivity Electrochemiluminescence-Based Detection System for Automated PCR Product Quantitation," <i>BioTechniques</i> 15:152-59 (1993) |
| 27 | Felgner <i>et al.</i> , "Lipofection: A highly efficient, lipid-mediated DNA-transfection procedure," <i>Proc. Natl. Acad. Sci. USA</i> 84:7413-17 (1987) |
| 28 | Hall, <i>et al.</i> , "Mapping Labeled Sites in <i>Escherichia coli</i> Ribosomal RNA: Distribution of Methyl Groups and Identification of a Photoaffinity-Labeled RNA Region Putatively at the Peptidyltransferase Center," <i>Biochemistry</i> 24:5702-5711 (1985) |
| 29 | Happ <i>et al.</i> , "New Approach to the Synthesis of 2'(3')-O-Aminoacyl Oligoribonucleotides," <i>J. Org. Chem.</i> 52:5387-91 (1987) |
| 30 | Hardesty <i>et al.</i> , "Extension and Folding of Nascent Peptides on Ribosomes." <i>The Translational Apparatus</i> , Nierhaus <i>et al.</i> ed: New York and London; Plenum Press. p.347-358 (1993) |
| 31 | Hardesty <i>et al.</i> , "Ribosome function determined by fluorescence," <i>Biochimie</i> 74:391-401 (1992) |
| 32 | Heckler <i>et al.</i> , "T4 RNA Ligase Mediated Preparation of Novel "Chemically Misacylated" tRNA ^{Phe} s," <i>Biochemistry</i> 23:1468-73 (1984) |
| 33 | Heckler <i>et al.</i> , "Preparation of 2'(3')-O-Acyl-pCpA Derivatives as Substrates for T4 RNA Ligase-Mediated "Chemical Aminoacylation," <i>Tetrahedron</i> 40:87-94 (1984) |
| 34 | Herman, <i>et al.</i> , "Affinity Chromatography of DNA Labeled with Chemically Cleavable Biotinylated Nucleotide Analogs," <i>Analytical Biochemistry</i> 156:48-55 (1986) |
| 35 | Hemmila, I.A., <i>Chemical Analysis "Applications of Fluorescence in Immunoassays"</i> , (Wiley&Sons 1991) pp.138-159 |
| 36 | Houlihan, <i>et al.</i> , "Nitrobenzyl Ester Chemistry for Polymer Processes Involving Chemical Amplification," <i>Macromolecules</i> 21:2001-2006 (1988) |
| 37 | Hudson, "Methodological Implications of Simultaneous Solid-Phases Peptide Synthesis. 1. Comparison of Different Coupling Procedures," <i>J. Org. Chem.</i> 53:617-624 (1988) |
| 38 | Johnson <i>et al.</i> , "Protein Synthesis and Secretion as seen by the Nascent Protein Chain," <i>The Translational Apparatus</i> , Nierhaus <i>et al.</i> ed: New York and London; Plenum Press. p. 359-370 (1993) |
| 39 | Johnson <i>et al.</i> , "N-Acetyllysine Transfer Ribonucleic Acid: A Biologically Active Analogue of Aminoacyl Transfer Ribonucleic Acids," <i>Biochemistry</i> 15:569-575 (1976) |
| 40 | Johnson, "Chemically Modified Aminoacyl-tRNA as a Probe of Ribosome Structure: the Synthesis and in vitro Activity of ϵ -N-acetyl-Lys-tRNA," 1973 Thesis Excerpts, University of Oregon, Eugene, Oregon |
| 41 | Kalbag and Roeske, "A Photolabile Protecting Group for Histidine," <i>J. Am. Chem. Soc.</i> 97:440-441 (1975) |
| 42 | Karolin, J., <i>et al.</i> , "Fluorescence and Absorption Spectroscopic Properties of Dipyrrometheneboron Difluoride (BODIPY) Derivatives in Liquids, lipid Membranes and Proteins," <i>J. Am. Chem. Soc.</i> 116:7801-7806 (1994) |
| 43 | Kozak, "Point Mutations Define a Sequence Flanking the AUG Initiator Codon that Modulates Translation by Eukaryotic Ribosomes," <i>Cell</i> 44:283-292 (1986) |
| 44 | Krafft, <i>et al.</i> , "Photoactivable Fluorophores. 3. Synthesis and Photoactivation of Fluorogenic Difunctionalized Fluoresceins," <i>J. Am. Chem. Soc.</i> 110:301-303 (1988) |
| 45 | Krieg <i>et al.</i> , "Photocrosslinking of the signal sequence of nascent preprolactin to the 54-kilodalton polypeptide of the signal recognition particle," <i>Proc. Natl. Acad. Sci. USA</i> 83:8604-08 (1986) |
| 46 | Kudlicki <i>et al.</i> , "Chaperone-dependent Folding and Activation of Ribosome-bound Nascent Rhodanese," <i>J. Mol. Biol.</i> 244:319-331 (1994) |
| 47 | Kurzchalia, T.V., <i>et al.</i> , "tRNA-Mediated Labelling of Proteins with Biotin. A Nonradioactive Method for the Detection of Cell-Free Translation Products," <i>Eur. J. Biochem.</i> 172:663-668 (1988) |
| 48 | Lesnikowski and Jaworska, "Studies on Stereospecific Formation of P-Chiral Internucleotide Linkage. Synthesis of (Rp,Rp)- and (Sp,Sp)-Thymidyl(3',5') Thymidyl(3',5') Thymidine Di(o,O-Phosphorothioate) using 2-Nitrobenzyl Group as a New S-Protection," <i>Tetrahedron Letters</i> 30(29):3821-3824 (1989) |
| 49 | Ma <i>et al.</i> , "In Vitro Protein Engineering Using Synthetic tRNA ^{Asp} with Different Anticodons," <i>Biochemistry</i> 32:7939-7945 (1993) |
| 50 | Markings and Tsien, "Caged Nitric Oxide, Stable organic Molecules from which Nitric Oxide can be Photoreleased," <i>J. Biological Chemistry</i> 269:6282-6285 (1994) |
| 51 | McCray <i>et al.</i> , "Properties and uses of photoreactive caged compounds," <i>Annu Rev Biophys Biophys Chem</i> 18:239-270 (1989) |

Examiner:

Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449 (Modified)		U.S. Department of Commerce Patent and Trademark Office		Attorney Docket No.: AMBER-06803	Serial No.: 10/034,736
				Applicant: Kenneth J. Rothschild <i>et al.</i>	
				Filing Date: 12/27/01	Group Art Unit: 1636
INFORMATION STATEMENT BY APPLICANT (Use Several Sheets If Necessary)					
(37 CFR § 1.98(b))					
OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)					
	52	Mendel, <i>et al.</i> , "Construction of a Light-Activated Protein by Unnatural Amino Acid Mutagenesis," <i>J. Am. Chem. Soc.</i> 113:2758-2760 (1991)			
	53	Miknis and Williams, "Total Synthesis of (±)-Aspirochlorine," <i>J. Am. Chem. Soc.</i> 115:536-547 (1993)			
	54	Molecular Cell Biology, J. Darnell <i>et al.</i> , Editors. Scientific American Books, N.Y., N.Y. pp.119-132 1991			
	55	Mouton, <i>et al.</i> , "A Reagent for Covalently Attaching Biotin to Proteins via a Cleavable Connector Arm," <i>Archives of Biochemistry and Biophysics</i> 218(1):101-108 (1982)			
	56	Neu and Heppel, "Nucleotide Sequence Analysis of Polyribonucleotides by Means of Periodate Oxidation Followed by Cleavage with an Amine," <i>J. Biol. Chem.</i> 239:2927-34 (1964)			
	57	Noren <i>et al.</i> , "A General Method for Site-Specific Incorporation of Unnatural Amino Acids into Proteins," <i>Science</i> 244:182-188 (1989)			
	58	Nishikubo, <i>et al.</i> , "Study of Photopolymers. XXXIV. Etherification and Esterification Reactions of Polymers with (o, m, or p)-Bromomethylnitrobenzene Using the DBU Method and the Photochemical Properties of the Resulting Polymers," <i>J. Polymer Science: Part A: Polymer Chemistry</i> 28:105-117 (1990)			
	59	Odom, <i>et al.</i> , <u>Methods in Molecular Biology</u> "In Vitro Engineering Using Acyl-Derivatized tRNAs," edited by: R. Martin Humana Press Inc., Totowa, NJ Chapter 7 pp. 93-103			
	60	Odom <i>et al.</i> , "Movement of tRNA but Not the Nascent Peptide during Peptide Bond Formation on Ribosomes," <i>Biochemistry</i> 29:10734-10744 (1990)			
	61	Oesterhelt <i>et al.</i> , "Bacteriorhodopsin: a biological material for information processing," <i>Quart. Rev. Biophys.</i> 4:425-78 (1991)			
	62	Ohtsuka, <i>et al.</i> , "Studies on Transfer Ribonucleic Acids and Related Compounds. 20. A New Versatile Ribooligonucleotide Block with 2'-(o-Nitrobenzyl) and 3'-Phosphorodanilidate Groups Suitable for Elongation of Chains in the 3' and 5' Directions," <i>J. Am. Chem. Soc.</i> 100(14):4580-4584 (1978)			
	63	Ohtsuka, <i>et al.</i> , "Studies on Transfer Ribonucleic Acids and Related Compounds; XVIII. A Photolabile 2'-Ether of Guanosine as an Intermediate for Oligonucleotide Synthesis," <i>Synthesis</i> 7:453-454 (1977)			
	64	Patchornik <i>et al.</i> , "Photosensitive Protecting Groups," <i>J. Am. Chem. Soc.</i> 92:6333-35 (1970)			
	65	Perri, <i>et al.</i> , "Tandem Photochemical Synthesis of N-Amino β-Lactams from Pyrazolidin-3-ones," <i>J. Org. Chem.</i> 55:6037-6047 (1990)			
	66	Peyser and Flechtner, "N-(α-Hydroxy-2-nitrosobenzyl)-1-naphthamide: A Photochemical Intermediate," <i>J. Org. Chem.</i> 25:4645-4646 (1987)			
	67	Picking <i>et al.</i> , "The use of synthetic tRNA as probes for examining nascent peptides on <i>Escherichia coli</i> ribosomes," <i>Biochimie</i> 73:1101-1107 (1991)			
	68	Picking <i>et al.</i> , "Evidence for RNA in the Peptidyl Transferase Center of <i>Escherichia coli</i> Ribosomes as Indicated by Fluorescence," <i>Biochemistry</i> 31:12565-12570 (1992)			
	69	Picking <i>et al.</i> , "The Conformation of Nascent Polylysine and Polyphenylalanine Peptides on Ribosomes," <i>J. of Biological Chemistry</i> 266:1534-1542 (1991)			
	70	Picking <i>et al.</i> , "Fluorescence Characterization of the Environment Encountered by Nascent Polyalanine and Polyserine as They Exit <i>Escherichia coli</i> Ribosomes during Translation," <i>Biochemistry</i> 31:2368-2375 (1992)			
	71	Picking <i>et al.</i> , "A synthetic alanyl-initiator tRNA with initiator tRNA properties as determined by fluorescence measurements: Comparison to a synthetic alanyl-elongator tRNA," <i>Nucleic Acids Research</i> 19:5749-5754 (1991)			
	72	Pillai, "Photoremovable Protecting Groups in Organic Synthesis," <i>Synthesis</i> 1-26 (1980)			
	73	Powell <i>et al.</i> , "Molecular Diagnosis of Familial Adenomatous Polyposis," <i>N. Engl. J. Med.</i> 329:1982-87 (1993)			
Examiner:			Date Considered:		
EXAMINER: Initial citation considered. Draw line-through citation if not-in conformance and not considered. Include copy of this form with next communication to applicant.					

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: AMBER-06803

Serial No.: 10/034,736

INFORMATION DISCLOSURE STATEMENT BY APPLICANT

(See Several Sheets If Necessary)

Applicant: Kenneth J. Rothschild *et al.*

Filing Date: 12/27/01

Group Art Unit: 1636

(37 CFR § 1.98(b))

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

74	Pratt, "Coupled Transcription-Translation in Prokaryotic Cell-Free System," (<i>Transcription and Translation</i> , B.D. Hames and S.J. Higgins, Editors, p. 179-209, IRL Press, Oxford, 1984)
75	Promega Technical Bulletin No. 182; tRNA ^{nucend} TM : Non-Radioactive Translation Detection System. 1-16, Sept. 1993
76	Renil and Pillai, "Synthesis of fully Protected Peptides on a Tetraethyleneglycol Diacrylate (TTEGDA)-Crosslinked Polystyrene support with a Photolytically Detachable 2-Nitrobenzyl Anchoring group," <i>Tetrahedron Letters</i> 35(22):3809-3812 (1994)
77	Robertson, <i>et al.</i> , "A General and Efficient Route for Chemical Aminoacylation of Transfer RNAs," <i>J. Am. Chem. Soc.</i> 113:2722-2729 (1991)
78	Robertson, <i>et al.</i> , "The use of 5'-phospho-2 deoxyribocytidylylriboadenosine as a facile route to chemical aminoacylation of tRNA," <i>Nucleic Acids Research</i> 17(23):9649-9660 (1989)
79	Sampson and Uhlenbeck, "Biochemical and physical characterization of an unmodified yeast phenylalanine transfer RNA transcribed <i>in vitro</i> ," <i>Proc. Natl. Acad. Sci. USA</i> 85:1033-37 (1988)
80	Seong and RajBhandary, "Escherichia coli formylmethionine tRNA: Mutations in ^{GGG} sequence conserved in anticodon stem of initiator tRNAs affect initiation of protein synthesis and conformation of anticodon loop," <i>Proc. Natl. Acad. Sci. USA</i> 84:334-338 (1987)
81	Shore <i>et al.</i> , "A Fluorescent Probe Capable of Incorporation into Nascent Polypeptide Chains," 1986 <i>Federation Proceedings</i> 45, 1566 Abstract
82	Shore <i>et al.</i> , "Accessibility of AA-tRNA and Nascent Chain During Protein Synthesis," 1988 <i>FASEB Journal</i> 2, A1045 Abstract
83	Shore, "The Use of Fluorescent-Labeled Amino Acids to Examine the Environment of Ribosome-Bound Nascent Polypeptide Chains," 1991 Dissertation, University of Oklahoma, Norman Oklahoma
84	Sonar, S., <i>et al.</i> , "Cell-Free Synthesis, Functional Refolding and Spectroscopic Characteristics of Bacteriorhodopsin, an Integral Membrane Protein," <i>Biochemistry</i> 32:13777-13781 (1993)
85	Spirin <i>et al.</i> , "A Continuous Cell-Free Translation System Capable of Producing Polypeptides in High Yield," <i>Sci.</i> 242:1162-64 (1988)
86	Stephen, "High-Resolution Preparative SDS-Polyacrylamide Gel Electrophoresis: Fluorescent Visualization and Electrophoretic Elution-Concentration of Protein Bands," <i>Anal. Biochem.</i> 65:369-79 (1975)
87	Thiele and Fahemholz, "Photocleavable Biotinylated Ligands for Affinity Chromatography," <i>Analytical Biochemistry</i> 218:330-337 (1994)
88	Thompson, <i>et al.</i> , "Photocleavable Nitrobenzyl-Protein Conjugates," <i>Biochemical and Biophysical Research Communications</i> 201(3):1213-1219 (1994)
89	Varshney, <i>et al.</i> , "Initiation of protein synthesis from a termination codon," <i>PNAS</i> 87:1586-1590 (1990)
90	Varshney, <i>et al.</i> , "Direct Analysis of Aminoacylation Levels of tRNAs <i>In Vivo</i> ," <i>J. Biological Chemistry</i> 266(36):24712-24718 (1991)
91	Whitney, "A Photochemical approach to the synthesis of (±)-biotin," <i>Can. J. Chem.</i> 59:2650-2653 (1981)
92	Widder <i>et al.</i> , "Magnetic Microspheres: A Model System for Site Specific Drug Delivery <i>In Vivo</i> ," <i>Proc. Soc. Exp. Biol. & Med.</i> 58:141-46 (1978)
93	Wilchek and Bayer, "Applications of Avidin-Biotin Technology: Literature Survey," <i>Methods in Enzymology</i> 184:14-45 (1990)
94	Yen, <i>et al.</i> , "Optically controlled ligand delivery, I," <i>Makromol. Chem.</i> 190:69-82 (1989)
95	Zehavi, <i>et al.</i> , "Light-Sensitive Glycosides. I. 6-Nitroveratryl β-D-Glucopyranoside and 2-Nitrobenzyl β-D-Glucopyranoside," <i>J. Org. Chem.</i> 37(14):2281-2285 (1972)
96	Zehavi and Herchman, "Enzymic synthesis of oligosaccharides on a polymer support, light-sensitive, water-soluble substituted poly(vinyl alcohol)," <i>Carbohydrate Research</i> 128:160-164 (1984)
97	Zehavi, <i>et al.</i> , "Enzymic Synthesis of Oligosaccharides on a Polymer Support Light-Sensitive, Substituted Polyacrylamide Beads," <i>Carbohydrate Research</i> 124:23-34 (1983)

Examiner:

Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.